Contents

Preface vii

Conference Organization viii

Word of Welcome ix

Yamada Science Foundation x

Executive Members of the Foundation xi

QUANTUM HALL STATES VERSUS ELECTRON SOLIDS

Optical spectroscopy of correlated phases of degenerate two-dimensional electrons
A.J. Turberfield, S.R. Haynes, P.A. Wright, R.A. Ford, R.G. Clark, J.F. Ryan, J.J. Harris and C.T. Foxon 1

Magneto-optics in the integer and fractional quantum Hall and electron solid regimes
B.B. Goldberg, D. Heiman, A. Pinczuk, L.N. Pfeiffer and K.W. West 9

Transport measurements on the electronic phases around $\nu = 1/5$ Landau level filling
H.W. Jiang, H.L. Stormer, D.C. Tsui, L.N. Pfeiffer and K.W. West 18

Wigner solid in random field: rigidity, pinning frequency and conduction threshold
F.I.B. Williams, G. Deville, D.C. Glattli, P.A. Wright, E.Y. Andrei, R.G. Clark, B. Etienne, C.T. Foxon and J.J. Harris 23

Magnetooptics of the incompressible Fermi liquid and of the Wigner solid
I.V. Kukushkin, N.J. Pulsford, K. von Klitzing, K. Ploog and V.B. Timofeev 30

Temperature dependence of the luminescence from a two-dimensional electron system in the Wigner solid regime
M. Hayne, M.K. Ellis, A.S. Plaut, A. Usher and K. Ploog 39

Studies of the reentrant insulating phase around the $\frac{1}{5}$ FQH liquid: I–V curves, noise, RF bias
L.W. Engel, T. Sajoto, Y.P. Li, D.C. Tsui and M. Shayegan 44

Transport properties of a 2D electron solid in Si-MOSFETs
M. D'Iorio, J.W. Campbell, V.M. Pudalov and S.G. Semenchinsky 49

Evidence for two-dimensional electron solid in Si inversion layers
S.V. Kravchenko, J.A.A.J. Perenboom and V.M. Pudalov 55

Behavior of electrons at soft edges in the fractional quantum Hall and Wigner crystal states
H.A. Fertig and S. Das Sarma 60

FRACTIONAL QUANTUM HALL EFFECT

Strongly correlated fractional quantum Hall states: description in terms of weakly interacting composite fermions
J.K. Jain 65

Quasiparticle charge of 1/3 FQH liquid from resistance fluctuation measurements
S.W. Hwang, J.A. Simmons, D.C. Tsui and M. Shayegan 72

Microwave response of quasi-particles in the FQHE
R. Meisels, F. Kuchar, J.J. Harris and C.T. Foxon 76

The fractional quantum Hall effect in high mobility two-dimensional hole gases
A.G. Davies, R. Newbury, M. Pepper, J.E.F. Frost, D.A. Ritchie and G.A.C. Jones 81

Fractional quantum Hall effect in thick layers and double quantum well systems
S. He, X.C. Xie and S. Das Sarma 87

INTEGER QUANTUM HALL EFFECT

Spatial potential distribution in GaAs/$Al_xGa_{1-x}As$ heterostructures under quantum Hall conditions studied with the linear electro-optic effect
P.F. Fontein, P. Hendriks, F.A.P. Blom, J.H. Wolter, L.J. Giling and C.W.J. Beenakker 91

Image of the dissipation in gated quantum Hall effect samples
U. Klass, W. Dietsche, K. von Klitzing and K. Ploog 97

Hall effect and localization in the Landau subbands of Si inversion layers
J. Wakabayashi 100

Direct measurement of critical exponents in the quantum Hall regime
S. Koch, R.J. Haug, K. von Klitzing and K. Ploog 108

A direct comparison of the quantized Hall resistance in high critical current gallium arsenide and silicon devices
A. Hartland, K. Jones, J.M. Williams, T. Galloway, B.L. Gallagher, C.R.H. White and M. Henini 112

Anomalously offset quantized Hall plateaus in high-mobility Si-MOSFETs
C.T. Van Degrift, K. Yoshihiro, M.E. Cage, D. Yu, K. Segawa, J. Kinoshita and T. Endo 116

Capacitively-coupled measurements on the frequency dependent diagonal magnetoconductance of a 2DEG in GaAs heterostructure in the quantum Hall regime
J.I. Lee, P.J. Stiles and M. Heiblum 120

Characterization of the electronic states near the centres of the Landau bands under quantum Hall conditions
B. Huckestein, B. Kramer and L. Schweitzer 125

Effect of probes in 2D electron systems with finite width subject to strong magnetic fields
Y. Ono 129

Electronic states in disordered layered systems in the quantum Hall regime
T. Ohtsuki, Y. Ono and B. Kramer 134

The quantum Hall effect in anomalous band structures
H. Aoki 137

TRANSPORT PROPERTIES

Interactions between hot injected electrons and the cold electrons in the two-dimensional electron gas base of a vertical hot electron transistor
P. Matthews, M.J. Kelly, D.G. Hasko, H. Ahmed, J.E.F. Frost, D.A. Ritchie and G.A.C. Jones 141

Magnetotransport in planar doping structures
X.-m. Feng, J.J. Mares, M.E. Raikh, F. Koch, D. Grützmacher and A. Kohl 147

Magnetostransport properties of double-layer systems in wide single quantum wells
Y.W. Suen, J. Jo, M.B. Santos, L.W. Engel and M. Shayegan 152

Magneto-transport study in restricted MOS inversion layers
A. Ohata and A. Toriumi 157

Conductance in ultra-short channel Si MOSFETs
A. Hartstein 162

A new method for measuring the effective diffusion constant of two-dimensional electrons in the presence of magnetic fields
H. Shao, H. Zheng and Y. Li 167

Superlattice Bloch electron conduction and negative differential mobility
X.L. Lei, N.J.M. Horing and H.L. Cui 172

Two-dimensional hole gas in Si/SiGe heterostructures
F.F. Fang, P.J. Wang, B.S. Meyerson, J.J. Nocera and K.E. Ismail 175

A transport model for a two-dimensional lateral-surface-superlattice
M.H. Degani and J.P. Leburton 179

180° phase shift of phonon drag magnetothermopower oscillations in high mobility 2DEGs
T.M. Fromhold, P.N. Butcher, G. Qin, B.G. Mulimani, J.P. Oxley and B.L. Gallagher 183

Frequency-dependent conductivity of a pinned Wigner crystal
R. Côté and A.H. MacDonald 187

Electron motion on a curved interface
M. Ikegami and Y. Nagaoka 193

TUNNELING

Resonant magnetotunneling spectroscopy: a direct probe of the complicated dispersion curves and negative mass behaviour of holes confined in a quantum well
L. Eaves, R.K. Hayden, M.L. Leadbeater, D.K. Maude, E.C. Valadares, M. Henini, F.W. Sheard, O.H. Hughes, J.C. Portal and L. Cury 199

Tunneling processes between quantum wires and two-dimensional electron states
J. Smoliner, W. Demmerle, F. Hirler, E. Gornik, G. Böhm and G. Weimann 207

Vertical transport in Schottky-gated, laterally confined double-barrier quantum well heterostructures
P. Guéret, N. Blanc, R. Germann and H. Rothuizen 212

Quantum transport phenomena in InAs/GaSb/InAs tunneling devices under high magnetic fields
T. Takamasu, N. Miura, K. Taira, K. Funato and H. Kawai 217

Tunneling spectroscopy of a 2DEG in InAs: experiment and theory
R. Winkler, U. Kunze and U. Rössler 222

Far infrared response of double barrier resonant tunneling structures
V.A. Chitta, R.E.M. de Bekker, J.C. Maan, S.J. Hawksworth, J.M. Chamberlain, M. Henini and G. Hill 227

Tunneling spectroscopy and miniband structure of selenium delta-doped GaAs
P. Basmaji, A.C. Notari, B. Schrappe, M.H. Degani, L. Ioriatti and O. Hipólito 231

Calculated contribution of localised LO phonon modes to the valley current of a double-barrier diode
A. Nogaret and J.C. Portal 234

BALLISTIC TRANSPORT

Ballistic electron optics
J. Spector, J.S. Weiner, H.L. Stormer, K.W. Baldwin, L.N. Pfeiffer and K.W. West 240

Classical and quantum transmission effects in submicron-size dots
R.P. Taylor, A.S. Sachrajda, J.A. Adams, P. Zawadzki, P.T. Coleridge and M. Davies 247

Ballistic electron propagation in macroscopic four-terminal structures and its gate controlled transmission and reflection
S. Tarucha, Y.K. Fukai, Y. Hirayama and T. Saku 253

Quantized thermoelectric power in the ballistic transport regime
Y. Okuyama, T. Sakuma and N. Tokuda 258

Quantized conductance of a quasi-ballistic one-dimensional wire
J. Wróbel, F. Kuchar, K.E. Ismail, K.Y. Lee, H. Nickel and W. Schlapp 261

Quantum oscillations of transverse resistance in series and single point-contact configurations
S. Yamada, M. Yamamoto and K. Aihara 265

EDGE STATES

Overshoot of quantum Hall plateaus
C.A. Richter, R.G. Wheeler and R.N. Sacks 270

Geometry-dependent resistances of two-dimensional electron gas at high magnetic fields
H. Nii, M. Ohsawa, S. Komiyama, S. Fukatsu, Y. Shiraki, R. Itoh and H. Toyoshima 275

Influence of interedge channel scattering on the magneto-transport of 2D-systems
G. Müller, E. Diessel, D. Weiss, K. von Klitzing, K. Ploog, H. Nickel, W. Schlapp and R. Lösch 280

Spin dependence of inter-edge-channel scattering in Si-MOSFETs in the quantum Hall regime
P.C. van Son, S.L. Wang and T.M. Klapwijk 284

Selective backscattering and the breakdown of the quantum Hall effect
L.W. Molenkamp, M.J.P. Brugmans, H. van Houten, C.W.J. Beenakker and C.T. Foxon 288

Influence of a constriction on the longitudinal resistance of a 2DEG: effects of electrical fields and edge states
G. Hein, P. Schneider, L. Schweitzer, F.J. Ahlers, L. Bliek, H. Nickel, R. Lösch and W. Schlapp 293

Non-local magnetoresistance at the crossover between the classical and quantum transport regimes
A.K. Geim(Heym), P.C. Main, P.H. Beton, P. Středa, L. Eaves, C.D.W. Wilkinson and S.P. Beaumont 298

Nonlocal Shubnikov–de Haas effect in Si-MOSFETs
K. Oto, S. Takaoka, K. Murase, S. Suyama and T. Serikawa 303

QUANTUM DOTS AND WIRES

Static and dynamic conductivity of lateral superlattices
A. Lorke 307

Magnetoresistance in a grid-type lateral superlattice: the role of disorder
D. Weiss, A. Menschig, K. von Klitzing and G. Weimann 314

Anisotropic electron transport through a rectangular antidot lattice
K. Ensslin, S. Sasa, T. Deruelle and P.M. Petroff 319

Magnetoresistance oscillations of a 2DEG in a weak lateral superlattice: manifestation of "Hofstadter's butterfly"
R.R. Gerhardts and D. Pfannkuche 324

Quantum Hall and transmission resonances of quantum dot arrays: a possible spectroscopy of Hofstadter butterflies
G. Kirczenow 330

Self-consistent calculations on confined electrons in three-dimensional geometries
A. Kumar 335

Direct magneto-optical observation of a quantum confined one-dimensional electron gas
A.S. Plaut, H. Lage, P. Grambow, D. Heitmann, K. von Klitzing and K. Ploog 341

Optical singularities of the one-dimensional electron gas in semiconductor quantum wires
J.M. Calleja, A.R. Goñi, B.S. Dennis, J.S. Weiner, A. Pinczuk, S. Schmitt-Rink, L.N. Pfeiffer, K.W. West, J.F. Müller and A.E. Ruckenstein 346

Plasma oscillations in quasi-classical dots
S. Nazin, K. Tevosyan and V. Shikin 351

Spectrum of collective modes and possible symmetry breaking transitions in arrays of quantum dots
A.V. Chaplik and L. Ioriatti 354

Far-infrared transmission spectra of laterally confined 2D plasma in quantum wires: influence of a magnetic field
S. Katayama 359

Many-particle effects in quantum well wires: plasmon–phonon coupling and depolarization effects
L. Wendler, R. Haupt and R. Pechstedt 363

Magnetotransport in Coulomb coupled quantum wires
P. Vasilopoulos and H.C. Tso 368

One-dimensional electron transport on the surface channel of InAs quantum wells
R.J. Haug, H. Munekata and L.L. Chang 374

Conductance fluctuations in a narrow GaAs/AlGaAs split gate wire
K. Ishibashi, Y. Aoyagi, S. Namba, Y. Ochiai, J.P. Bird and M. Kawabe 378

Conductance fluctuations in mesoscopic quantum wires near the ballistic regime
H. Higurashi, S. Iwabuchi and Y. Nagaoka 382

Size effect in the low temperature magnetoresistance of narrow GaAs/AlGaAs wires
Y. Ochiai, T. Onishi, J.P. Bird, M. Kawabe, K. Ishibashi, Y. Aoyagi and S. Namba 388

Aharonov–Bohm effect under high magnetic field in a Corbino disk anti-dot channel
K. Tsubaki, T. Honda and Y. Tokura 392

COULOMB BLOCKADE

The frequency-locked turnstile device for single electrons
L.J. Geerligs 396

Quantized current in a quantum dot turnstile
L.P. Kouwenhoven, A.T. Johnson, N.C. van der Vaart, D.J. Maas, C.J.P.M. Harmans and C.T. Foxon 405

Variable range hopping via a small number of localized states in nanometer scale MOSFETs
C. de Graaf, J.W.G. Wildöer, J. Caro, S. Radelaar and K. Heyers 409

Electron transport through one quantum dot and through a string of quantum dots
R.J. Haug, J.M. Hong and K.Y. Lee 415

Coulomb blockade of tunneling in a 2D electron gas
C. Pasquier, D.C. Glattli, U. Meirav, F.I.B. Williams, Y. Bin and B. Etienne 419

Scattering and Coulomb blockade in magnetotunneling across singly and multiply connected barriers in quasi-two-dimensional systems
C. Tejedor and J.J. Palacios 424

Gate-controlled conductance oscillations in small one-dimensional electron cavities
F.M. de Aguiar, D.A. Wharam, T. Heinzel, A. Lorke and J.P. Kotthaus 428

Charging energy and collective response of a quantum dot resonant tunneling device
M.P. Stopa 433

Single electron tunnelling through a donor state in a gated resonant tunnelling device
M.W. Dellow, C.J.G.M. Langerak, P.H. Beton, T.J. Foster, P.C. Main, L. Eaves, M. Henini, S.P. Beaumont and C.D.W. Wilkinson 438

Coulomb blockade oscillations in semiconductor nanostructures
H. van Houten 442

MANY-BODY EFFECTS

Electron–electron scattering between parallel two-dimensional electron gases
T.J. Gramila, J.P. Eisenstein, A.H. MacDonald, L.N. Pfeiffer and K.W. West 446

Magneto-rotons in wide parabolic quantum wells
K. Karraï, X. Ying, H.D. Drew, M.B. Santos, M. Shayegan, S.-R.E. Yang and A.H. MacDonald 451

Many-body effects in dense electron–hole plasma in InGaAs QWs with simple and complex valence band
L.V. Butov, V.D. Kulakovskii, T.G. Andersson, A. Forchel and D. Grützmacher 457

Screening and correlation effects in degenerately center doped GaAs/AlGaAs single quantum wells
C.I. Harris, H. Kalt, B. Monemar and K. Köhler 462

Single-particle and many-body effects in the edge magnetoplasma excitations of a two-dimensional electron system
I. Grodnensky, D. Heitmann and K. von Klitzing 467

Exchange and correlation in quasi-two-dimensional plasmas in quantum well structures
D. Gammon, B.V. Shanabrook, J.C. Ryan, G. Brozak and D.S. Katzer 471

Plasmon excitation by dual optical couplers
R.J. Wilkinson, C.D. Ager and H.P. Hughes 475

Magnetic-field-induced excitonic instability in the 2D electron gas
G.E.W. Bauer 482

EXCITONS

Two-dimensional exciton magnetic polaron in semimagnetic quantum wells
D.R. Yakovlev, I.N. Uraltsev, W. Ossau, G. Landwehr, R.N. Bicknell-Tassius, A. Waag and S. Schmeusser 485

Enhanced radiative recombination of free excitons in GaAs quantum wells
B. Deveaud, F. Clérot, N. Roy, B. Sermage and D.S. Katzer 491

Indirect excitons in coupled quantum well structures
A. Zrenner, P. Leeb, J. Schäfer, G. Böhm, G. Weimann, J.M. Worlock, L.T. Florez and J.P. Harbison 496

Thermodynamics of excitons in coupled quantum wells
J.A. Kash, M. Zachau, E.E. Mendez, J.M. Hong and T. Fukuzawa 502

Excitonic absorption spectra and Sommerfeld factors of one-dimensional systems
T. Ogawa and T. Takagahara 506

Two-photon absorption spectra of low-dimensional semiconductors
A. Shimizu, T. Ogawa and H. Sakaki 512

OPTICAL PROPERTIES

Infrared spectroscopy of subbands, minibands, and donors in GaAs/AlGaAs superlattices
M. Helm, F.M. Peeters, F. DeRosa, E. Colas, J.P. Harbison and L.T. Florez 518

Electronic Raman scattering on highly doped single quantum wells
B. Jusserand, D. Richards, B. Etienne, H. Peric and G. Fasol 527

Multiply resonant Raman scattering in Stark ladder superlattices
H. Schneider, J. Wagner, K. Ploog and K. Fujiwara 531

Spatially direct and indirect optical transitions in shallow etched GaAs/AlGaAs quantum wires
F. Hirler, R. Küchler, R. Strenz, G. Abstreiter, G. Böhm, J. Smoliner, G. Tränkle and G. Weimann 536

Temperature dependent electronic properties of HgZnTe/CdTe superlattices
Y. Guldner, J. Manassès, J.P. Vieren, M. Voos and J.P. Faurie 541

Magnetic field induced metal–insulator transitions in modulation-doped multiple-quantum-well structures
Y.J. Wang, B.D. McCombe and W. Schaff 547

Magnetic field induced indirect gap in a modulation doped quantum well
D.M. Whittaker, T.A. Fisher, P.E. Simmonds, M.S. Skolnick, R.S. Smith, L.L. Taylor and S.J. Bass 549

Kinetics of pressure induced Γ–χ electron transfer in a type-I superlattice
J. Nunnenkamp, K. Reimann, J. Kuhl and K. Ploog 553

Effects of localization on optical spectra for shallow acceptors in center-doped GaAs/AlGaAs multiple quantum wells
B. Monemar, P.O. Holtz, P. Bergman, C.I. Harris, H. Kalt, M. Sundaram, J.L. Merz and A.C. Gossard 556

Photoinduced intersubband absorption in n-type well- and barrier-doped quantum wells
Y. Garini, E. Cohen, E. Ehrenfreund, A. Ron, K.K. Law, J.L. Merz and A.C. Gossard 561

Electronic properties of two-dimensional D^- centres
S. Huant, A. Mandray, G. Martinez, M. Grynberg and B. Etienne 565

Temperature dependence of the magneto-optical properties of a $CdTe/Cd_{1-x}Mn_xTe$ multiquantum well structure
E. Deleporte, G. Peter, J.M. Berroir, C. Delalande, J.M. Hong and L.L. Chang 570

Piezoelectric control of doping and band structure in the crossed gap system GaSb/InAs
M. Lakrimi, C. López, R.W. Martin, G.M. Summers, G.M. Sundaram, K.S.H. Dalton, N.J. Mason, R.J. Nicholas and P.J. Walker 575

Utilization of a piezoelectric field in a one-sided-doped quantum well
U. Ekenberg and D. Richards 580

Resonant decay of pair excitations of a 2D hole gas into optical phonons
F. Bechstedt and H. Gerecke 585

Filling factor dependent Landau level broadening studied with inter- and intraband magneto-optics in GaInAs/AlInAs MDQWs
M. Potemski, Y.H. Zhang, M. Hausenblas, J.C. Maan, G. Martinez and K. Ploog 591

Properties of 2D electrons and associated bound donors in delta-doped InSb
M.J. Yang, W.J. Moore, R.J. Wagner, J.R. Waterman, C.H. Yang, P.E. Thompson and J.L. Davis 595

Magneto-optical investigation of energy band structure in $GaAs/Al_xGa_{1-x}As$ short period superlattices under pulsed high magnetic fields
S. Sasaki and N. Miura 600

Experimental determination of effective mass within the energy gap
Y. Murayama and J.-i. Kasai 604

The electronic structures of a GaAs/AlAs Fibonacci superlattice
K. Hirose and H. Kamimura 609

LUMINESCENCE

Luminescence from the degenerate two-dimensional electrons at an ultrahigh mobility heterojunction
S.R. Haynes, R.A. Ford, A.J. Turberfield, P.A. Wright, R.G. Clark, J.F. Ryan, J.J. Harris and C.T. Foxon 614

Effect of hole relaxation and uniaxial stress on luminescence polarization in quantum wells
T. Uenoyama and L.J. Sham 618

Luminescence studies from wires of varying aspect ratio in GaAs/GaAlAs quantum wells
W.E. Leitch, C.M. Sotomayor Torres, D. Lootens, S. Thoms, P. van Daele, C.R. Stanley, P. Demeester and S.P. Beaumont 622

Fermi edge singularity and screening effects in the luminescence spectra of Si or Be δ-doped GaAs
J. Wagner, P. Ganser, K. Köhler, A. Fischer and K. Ploog 628

Fermi-edge singularities as probes of a two-dimensional electron gas in GaAs heterostructures
M. Fritze, W. Chen, W. Wałecki, A.V. Nurmikko, J.M. Hong and L.L. Chang 633

Time-resolved optical recombination in the quantum Hall regime
M. Dahl, D. Heiman, A. Pinczuk, B.B. Goldberg, L.N. Pfeiffer and K.W. West 638

Observation of resonant electron capture in AlGaAs/GaAs quantum well structures
A. Fujiwara, S. Fukatsu, Y. Shiraki and R. Itoh 642

Many body resonant polaron interaction in quantum wells containing high densities of free carriers
P.E. Simmonds, M.S. Skolnick, T.A. Fisher, K.J. Nash and R.S. Smith 646

CYCLOTRON RESONANCE

Screening effects in the cyclotron resonance of 2D electrons in GaAs/AlGaAs heterostructures and single quantum wells
M. Besson, E. Gornik, G. Böhm and G. Weimann 650

Resonant magnetopolaron coupling to both polar and neutral optical phonons in the layer compound InSe
F.M. Peeters, X.-G. Wu, J.T. Devreese, M. Watts, R.J. Nicholas, D.F. Howell, L. van Bockstal, F. Herlach, C.J.G.M. Langerak, J. Singleton and A. Chevy 654

Photon-drag current in 2DEG in the vicinity of cyclotron resonance and its first subharmonic
A.P. Dmitriev, S.A. Emelyanov, S.V. Ivanov, P.S. Kop'ev, Ya.V. Terent'ev and I.D. Yaroshetsky 659

ELECTRONS ON LIQUID HELIUM

The conductivity and mobility of 2D nondegenerate electrons in the strong localization regime
P.W. Adams 663

Low-temperature features of quantum magnetoconductance of a two-dimensional electron gas on a liquid helium surface
Yu.P. Monarkha 668

Two-ripplon processes of electrons on the liquid ^{4}He surface
T. Kawaguchi, T. Ohtsuki and M. Saitoh 671

Temperature dependent microwave absorption of electrons on helium films
B. Lehndorff, C. Vossloh, T. Günzler and K. Dransfeld 674

A new low frequency propagating mode in a 2DEG in a magnetic field
M.J. Lea, A.O. Stone, P. Fozooni, P.J.M. Peters, A.M.L. Janssen and R.W. van der Heijden 677

Effect of quenched disorder on a two-dimensional classical Wigner crystal
A.B. Dzyubenko and Y.E. Lozovik 680

Author index 686

Subject index 702